



**INVESTIGATION OF HISTOLOGICAL EFFECTS OF IRON-
FORTIFIED RICE TO THE SELECTED ORGANS OF
Mus musculus (ALBINO MICE)**

**An Undergraduate Research Study presented to
The Biological Sciences Department
De La Salle University – Dasmariñas
College of Science and Computer Studies**

**In Partial Fulfillment of the Requirements for the degree
Bachelor of Science in Biology major in Human Biology**

Buenafe, John Dominique A.

Cortez, Lakan S.

HUB 31

March 2014



Acknowledgment

This research paper is made possible through the help and encouragement of the researchers' families, friends, classmates, and our outmost gratitude to the Almighty God.

The researchers would also like to thank Dr. Chester Saldaña, for supervising the experimental procedure, Ms. Hazel Anne L. Tabo, the adviser of this study for her guidance, avid checking and moral support in the study, our panels, Dr. Arnold Fonollera, Dr. Rubie Causaren, and Dr. Janet Macawile for their constructive criticisms. Also Ms. Cherry Cuevas, our department chair and Dr. Johnny Ching, the Dean of the college.

Lastly, the researchers would like to thank their parents and guardians, Mrs. Zelda Soriano, Mr. Bonifacio. A. Buenafe, Mrs. Emerita A. Buenafe for their financial and moral support and encouragement. Also, Ms. Sarah Jane T. Canuto for prodding the researchers to pursue the study to the fullest.



Table of contents

ABSTRACT4

INTRODUCTION

Background of the Study.....5

Conceptual Framework7

Statement of the problem.....7

Hypotheses.....8

Scope and delimitation.....8

Significance of the study.....9

Definition of Terms.....10

REVIEW OF LITERATURE.....12

METHODOLOGY.....23

Research Design.....23

Research Setting.....23

Research Procedure.....24

Data gathering.....26

RESULTS AND DISCUSSION.....27

CONCLUSIONS AND RECOMMENDATIONS.....40



ABSTRACT

The study determined the histological effect of Iron-fortified rice diet on the selected organs of *Mus musculus* (albino mice) in comparison with normal rice diet. A total of 20 mice were divided equally into four different treatments for eight weeks. The control (T0) was given normal mice feed, treatment 1 (T1) was given mice feed mixed with iron-fortified rice with a ratio of 1:4, treatment 2 (T2) were given mice feed mixed with normal rice with the ration of 1:4, and treatment 3 (T3) was given normal rice mixed with iron-fortified rice with a ratio of 1:1 and the selected organs (pancreas, liver, spleen, brain, heart, and kidneys) of each albino mice were sampled for histological examination. The mice under T0 produced less evidence of histological change compared to T1, T2, T3, degrees of histological change for each treatment group in that order. There was higher incidence of hemosiderin deposits in T2 and T3 in which the researchers hypothesized that normal rice has an inhibitory effect on iron metabolism. The results highly suggest that iron-fortified rice in the diet of albino mice has little to no histologic effect on the selected organs compared to normal rice.