



**ANTIUROLITHIATIC POTENTIAL OF *Eleusine indica* L. (GOOSE GRASS) ROOT EXTRACT ON ETHYLENE GLYCOL INDUCED NEPHROLITHIASIS IN *Rattus norvegicus* (ALBINO RATS)**

A Research Presented to the  
Biological Sciences Department  
College of Science and Computer Studies  
De La Salle University-Dasmariñas  
City of Dasmariñas, Cavite

In Partial Fulfilment of the Requirements for the Degree  
Bachelor of Science in Medical Biology

**DANIEL OPOKU AMOAH**  
**MARITAN JOSON**

May 2017



### ABSTRACT

The study aimed to determine the antiurolithiathic potential of different concentrations *E. indica* root extract on ethylene glycol induced nephrolithiasis in *Rattus norvergicus* species of albino rats. It specifically targeted the effects of the plant extract on serum parameters (creatinine, BUN and uric acid) nitrituria, proteinuria, calcium oxalate excretion, and the histopathology of kidney tubules and glomeruli of albino rats. In the study, root of *E. indica* was collected, dried and extracted using distilled water of 80°C -100°C. The study involved 5 test groups; negative control group(T-), positive control group or lithogenic group (T+), two prophylactic groups (T<sub>1</sub> and T<sub>2</sub>) receiving different concentrations of *E. indica* root extract(T<sub>1</sub>- 500mg/kg and T<sub>2</sub>- 800mg/kg ), and one curative group receiving 800mg/kg of *E. indica* root extract. All the test groups received 0.75% ethylene glycol in water for 4 week ad libitum with the exception of the negative control. At the end of the study, there were elevated levels of serum creatinine, BUN and uric acid among the lithogenic group which was normalized in all the three treatment groups. There was no significant difference between the serum parameters of the *E. indica* treatment groups and that of negative control (P< 0.05). There was the presence of nitrituria, proteinuria, high pH and high levels of calcium oxalate excretion in the lithogenic group. But all these parameters were normalized in the three *E. indica* treatment groups. *E. indica* also maintained the normal structure of kidney glomeruli and tubules among T<sub>1</sub> and T<sub>2</sub> (prophylactic groups) and repaired kidney defects in T<sub>3</sub> (curative group). *E. indica* root extract thus possess antiurolithiathic potentials and can be used to prevent and cure nephrolithiasis in albino rats.

*Key Terms:* Nephrolithiasis, nitrituria, proteinuria, creatinine, ethylene glycol.



## TABLE OF CONTENTS

Title Page	1
Abstract	2
Approval sheet	3
Acknowledgments	4
Table of Contents	5
<b>CHAPTER 1 INTRODUCTION</b>	
1.1 Background of the Study	7
1.2 Conceptual Framework	9
1.3 Statement of the Problem	10
1.4 Scope and Limitations	11
1.5 Significance of the Study	12
1.6 Definition of Terms	13
<b>CHAPTER 2 LITERATURE REVIEW</b>	
2.1 Conceptual Literature	15
2.2 Related Studies	23
<b>CHAPTER 3 METHODOLOGY</b>	
3.1 Research Design	33
3.2 Research Procedure	34
3.3 Data Gathering and Statistical Analysis	37
<b>CHAPTER 4 RESULTS AND DISCUSSION</b>	
4.1 Results	39



4.2	Discussion	45
CHAPTER 5 CONCLUSION AND RECOMMENDATIONS		
5.1	Conclusion	56
5.2	Recommendations	57
CITED REFERENCES		58
APPENDICES		
A.	Photo Documentation	69
B.	Histopathology Plates	72
C.	Statistical Analysis	77
D.	Certifications	81
CURRICULUM VITAE		82