



**ANGIOGENIC EFFECTS OF *Spondias purpurea* L. (SINEGUELAS) LEAF
CRUDE EXTRACT ON THE CHORIOALLANTOIC MEMBRANE
OF A 12-DAY OLD DUCK EMBRYO**

An Undergraduate Research Presented to the
Biological Sciences Department
College of Science and Computer Studies
De La Salle University-Dasmariñas
City of Dasmariñas, Cavite

In Partial Fulfilment of the Requirements for the Degree
Bachelor of Science in Biology Major in Human Biology

LORRAINE E. CANALES
MIGUEL ANTONIO S. POBLETE

JUNE 2018



TABLE OF CONTENTS

Title Page	1
Approval Sheet	2
Acknowledgments	3
Table of Contents	5
Abstract	7
CHAPTER 1 INTRODUCTION	
1.1 Background of the Study	8
1.2 Conceptual Framework	10
1.3 Statement of the Problem	11
1.4 Hypotheses	11
1.5 Scope and Limitations	12
1.6 Significance of the Study	13
1.7 Definition of Terms	14
CHAPTER 2 LITERATURE REVIEW	
2.1 Conceptual Literature	15
2.2 Related Studies	19
CHAPTER 3 METHODOLOGY	
3.1 Research Design	24
3.2 Research Setting	24
3.3 Research Procedure	24



3.4	Data Gathering and Statistical Analysis	26
CHAPTER 4 RESULTS AND DISCUSSION		
4.1	Results	28
4.2	Discussion	30
CHAPTER 5 SUMMARY, CONCLUSION AND RECOMMENDATIONS		
5.1	Summary	34
5.2	Conclusion	34
5.3	Recommendations	35
	(Cited) References	37
Appendices		
A.	Photo Documentation	46
B.	Standard Procedure	47
C.	Photo Documentation	53
D.	Computation	57
E.	Raw Data	59
F.	Average Number of Collaterals	61
G.	Phytochemical Results	62
H.	One-Way Anova Results	63
	Curriculum Vitae	64



ABSTRACT

Angiogenesis is a physiological process for the formation of new blood vessels, and is essential in bringing nutrients and oxygen to a malignant tumor whose development in growth is not contained. The study tested the angiogenic potential of *Spondias purpurea* (sineguelas) to the chorioallantoic membrane of a 12-day old duck embryo. The leaf samples were collected and macerated for 48 hours. The mixture that it produced was filtered using a cheesecloth, and its filtrate was concentrated using a rotary evaporator. Various concentrations of 100 ppm, 200 ppm, and 300 ppm of each leaf crude extract were prepared and administered to the CAM of the duck embryo. After two days of incubation, the eggs were observed under a dissecting microscope and the number of collaterals were then counted. Based on the statistical analysis (Kruskal-Wallis Test), the control group, 100 ppm, 200 ppm, and 300 ppm of sineguelas crude leaf had no significant difference. This may be due to the phytochemicals present in the sineguelas, such as: flavonoid and phenols which can affect various steps in angiogenesis.

Keywords: angiogenesis, chorioallantoic membrane, Spondias purpurea, 12-day old duck egg