DIVERSITY OF DIATOMS IN RELATION TO THE PHYSICO-CHEMICAL CHARACTERISTICS OF DASMARINAS RIVER, DASMARIÑAS CAVITE

An Undergraduate Research Presented to The Biological Science Department College of Science De La Salle University – Dasmarinas

In Partial Fulfillment Of the Requirements for the Degree Bachelor of Science in Biology Major in Human Biology

Casareo, Norrie – Anne A. Blasquino, Alvin Dale Roel V.

Adviser: Ms. Ma. Luisa Cuaresma, MSc.

March 2009

TABLE OF CONTENTS

TITLE PAGE .		1
APPROVAL SH	IEET	2
ABSTRACT .		10
ACKNOWLED	GEMENT	11
CHAPTERS		
I. INTRO	DUCTION	
1	.1 Background of the Study	12
1	.2 Theoretical Framework	13
	.3. Statement of the Problem	14
di la constante	.4 Scope and Delimitation	14
1	.5 Significance of the Study	15
1	.6 Definition of Terms	16
II. REVI	EW OF RELATED LITERATURE	
2	.1 Conceptual Literature	17
2	2.2 Related Literature	23
III. MET	THODOLOGY	
3	.1 Research Design	27
3	2.2 Research Setting	28
3	.3 Research Procedure	28
3	.4 Gathering of Data and Statistical Treatment	30

IV. RESULTS AND DISCUSSION

4.1 Results
4.2 Discussion
V. CONCLUSIONS AND RECOMMENDATIONS
5.1 Conclusion
5.2 Recommendation
LITERATURE CITATION 54
APPENDICES
PHOTODOCUMENTATION
CERTIFICATION FROM NATONAL MUSEUM 104
GANTT CHART 105
BUDGET PROPOSAL 106
CURRICULUM VITAE

Since 1077 *

LIST OF TABLES

TABLE 4.1: Diatom Species collected	38
TABLE 4.2: Physico-chemical Characteristics of the river	41
TABLE 4.3: Equivalent diversity index range	42
TABLE 4.4: Simpson and Shannon diversity index, dry season	42
TABLE 4.5: Simpson and Shannon diversity index, rainy season	42
TABLE 4.6: Correlation of the Diversity of Diatoms, dry season	45
TABLE 4.7: Correlation of the Diversity of Diatoms, rainy season	46
TABLE 1: The collection of samples	64
TABLE 2: Salinity data of Dasmarinas River	65
TABLE 3: Total Dissolved Solids data of Dasmarinas River	65
TABLE 4: Dissolved Oxygen data of Dasmarinas River	65
TABLE 5: Biochemical oxygen demand data of Dasmarinas River	66
TABLE 6: Temperature data of Dasmarinas River	66
TABLE 7: pH data of Dasmarinas River	66
TABLE 8: Transparency data of Dasmarinas River	67
TABLE 9: Diatom Species, Dry season Upstream	68
TABLE 10: Diatom species, Dry season Midstream	69
TABLE 11: Diatom Species, Dry season Downstream	70
TABLE 12: Diatom Species, Rainy season Upstream	71
TABLE 13: Diatom Species, Rainy season Midstream	72

TABLE14: Diatom species, Rainy season, Downstream		73
TABLE 15: Diversity Indices, Upstream 1 of Dry season		74
TABLE 16: Diversity Indices, Upstream 2 of Dry season		74
TABLE 17: Diversity Indices, Upstream 3 of Dry season		75
TABLE 18: Diversity Indices, Midstrean 1 of Dry season		75
TABLE 19: Diversity Indices, Midstream 2 of Dry season		76
TABLE 20: Diversity Indices, Midstream 3 of Dry season		76
TABLE 21: Diversity Indices, Downstream 1 of Dry season		77
TABLE 22: Diversity Indices, Downstream 2 of Dry season		77
TABLE 23: Diversity Indices, Downstream 3 of Dry season		78
TABLE 24: Diversity Indices, Upstream 1 of Rainy season		78
TABLE 25: Diversity Indices, Upstream 2 of Rainy season		79
TABLE 26: Diversity Indices, Upstream 3 of Rainy season	•••••	80
TABLE 27: Diversity Indices, Midstrean 1 of Rainy season		81
TABLE 28: Diversity Indices, Midstream 2 of Rainy season		82
TABLE 29: Diversity Indices, Midstream 3 of Rainy season		83
TABLE 30: Diversity Indices, Downstream 1 of Rainy season		84
TABLE 31: Diversity Indices, Downstream 2 of Rainy season	•••••	85
TABLE 32: Diversity Indices, Downstream 3 of Rainy season		86
TABLE 33: Correlation of the diversity of diatoms, dry season		87
TABLE 34: Correlation of the diversity of diatoms, rainy season		88

LIST OF FIGURES

FIGURE 4.1: Graph of the relative abundance during dry season	39
FIGURE 4.2: Graph of the relative abundance, rainy season	40
FIGURE 1: Map of the water system of Dasmariñas Cavite with Imus Rive	er 89
FIGURE 2: Map of the Water System of Dasmariñas Cavite	90
FIGURE 3: Overview of Dasmarinas River, Dasmarinas Cavite	91
FIGURE 4: Upstream	91
FIGURE 5: Midstream	92
FIGURE 6: Downstream	92
FIGURE 7: Achnanthes lanceolata	93
FIGURE 8: Coscinodiscus lacustris	93
FIGURE 9: Cyclotella kutzingiana	94
FIGURE 10: Cyclotella maneghiniana	94
FIGURE 11: Diatoma elongatum (colony)	95
FIGURE 12: Diatoma valgare	95
FIGURE 13: Melosira agussizii	96
FIGURE 14: Melosira islandica	96
FIGURE 15: Navicula gastrum	97
FIGURE 16: Neidium affine	97
FIGURE 17: Nitzschia nyassensis	98
FIGURE 18: Nitzschia subrostrata	98

FIGURE 19: Pinnularia gibba		99
FIGURE 20: Stauroneis anceps		99
FIGURE 21: Stephanodiscus hantzse	chii (with filtrates and without filtrates)	100
FIGURE 22: Surrirella elegans		100
FIGURE 23: Synedra acus		101
FIGURE 24: Synedra tabulate		101
FIGURE 25: Measuring of the chem	ical parameters of the river on field	. 102
FIGURE 26: Measuring of the transp	parency of the river, midstream	102
FIGURE 27: Cleaning of diatoms		103



LIST OF APPENDICES

Appendix A: Procedures	56
Appendix B: Identification of Diatoms by Sir Gapas	62
Appendix C: Sample Collection	64
Appendix D: Physico-chemical Caracteristics	65
Appendix E: Diatoms Sp. Relative Abundance	68
Appendix F: Diversity Index	74
Appendix G: Pearson Correlation	87
Appendix H: Maps	89
Appendix I: Dasmarinas River	91
Appendix J: Diatoms	93

ABSTRACT

The study entitled: Diversity of Diatoms in Relation to the Physico-chemical Characteristics of Dasmarinas River talks about the diatom species that can be found in Dasmarinas River in relation to its physico-chemical characteristics namely: pH, salinity, transparency, DO, temperature, TDS and BOD. Dasmarinas River is classified according to the water standard set by the DENR under Administrative Order No. 34 Series of 1990 and it is categorized under Class D which means that the river is suitable for use in irrigation, livestock watering and agriculture. Collections of samples were done twice, one during the dry season and the other during the rainy season from the duration of May to August. The species collection was done using plankton net with a mesh size of 2µm and was subjected to centrifugation using a rocket type centrifuge to isolate and identify the diatom species. Testing for its physico-chemical characteristics on the other hand was done on field except for the BOD. It was found out that more diatom species emerged during the rainy season because of factors like the fulvic acid content of the river and the richness of the environment after the rain. There are nineteen species of diatoms identified and were computed for its Simpson and Shannon indices, both used in identifying the diversity of diatoms. Their diversity was also correlated to the physico-chemical characteristics of the river using the Pearson r correlation and showed that the higher the diversity index of the species, the more diverse the organisms are. However, there are no rare species found in the river thus, the site might be ideal for industrial building purposes.