ANTIBACTERIAL ACTIVITY OF Cucurbita maxima (SQUASH) AND Citrullus lanatus (WATERMELON) SEED EXTRACTS AGAINST SELECTED GASTRO-INTESTINAL BACTERIAL PATHOGENS

A Research Presented to the
Biological Sciences Department
College of Science and Computer Studies
De La Salle University - Dasmariñas
City of Dasmariñas, Cavite

In Partial Fulfilment of the Requirements for the degree of Bachelor of Science in Biology Major in Human Biology

RAY AMELEE T. ENDAYA
INAH PATRICIA H. SUMAGUE

March 2014

ABSTRACT

The inhibitory activity of ethanolic seed extract of Citrullus lanatus (Watermelon) and Cucurbita maxima (squash) against selected gram-negative enteric pathogens, Salmonella typhi, Salmonella non-typhi, Vibrio choleare, and Shigella sp was evaluated using disk diffusion assay. Various concentrations of extracts (50%, 75% and 100%) were utilized as treatments. Result revealed that all treatments do not manifest inhibitory activity against the said gastrointestinal causing bacterial pathogens as compared to the commercially available antibiotic, Ciprofloxacin, that exhibit ability to destroy the integrity of bacterial DNA.

TABLE OF CONTENTS

Title Page	1			
Abstract	2			
Approval sheet	3			
Acknowledgements	4			
Table of Contents	5			
List of Tables	7			
List of Plates	8			
CHAPTER 1 INTRODUCTION				
1.1 Background of the Study	9			
1.2 Conceptual Framework	11			
1.3 Statement of the Problem	11			
1.4 Hypotheses	12			
1.5 Scope and Limitations	12			
1.6 Significance of the Study	13			
1.7 Definition of Terms	14			
CHAPTER 2 LITERATURE REVIEW				
2.1 Conceptual Literature	15			
2.2 Related Studies	21			
CHAPTER 3 METHODOLOGY				

24

24

25

28

29

35

48



Data Gathering and Statistical Analysis

CHAPTER 4 RESULTS AND DISCUSSION

4.2 Discussion	32

CHAPTER 5 CONCLUSION AND RECOMMENDATION

5.2 Recommendation	35
Cited References	36

Appendices

3.4

4.1 Results

5.1 Conclusion

D. Curriculum Vitae

A.	Taxonomic account		4′
В.	Standard procedure	office Con	42
C.	Photodocumentation	* Since 1977 • Dasmar	44

LIST OF TABLES

- Table 2.1 Phytochemical Components of C. maxima
- Table 2.1 Phytochemical Components of C. maxima
- Table 4.1. Zone of inhibition (mm) of *Cucurbita maxima* (squash) and *Citrullus lanatus* (watermelon) seed extracts against 10⁸cfu *V. cholera, Shigella sp.Salmonella* non-typhi and *Salmonella typhi*



LIST OF PLATES

- Plate 4.1 Inhibitory activity of *C. maxima* against 10⁸ cfu (A) *S. non-typhi,* (B) *S. typhi.* (C) *V. cholera, and* (D) *Shigella sp.*
- Plate 4.1 Inhibitory activity of *C. lanatus* against 10⁸ cfu (A) *S. non-typhi,* (B) *S. typhi.* (C) *V. cholera, and* (D) *Shigella sp.*

