CORRELATION BETWEEN THE AMOUNTS OF LEAD INCLAMS, MARINE WATER, AND BENTHIC SEDIMENTS OF ANG PULO MANGROVE RESERVE IN CALATAGAN, BATANGAS

An Undergraduate Research Presented to

The Faculty of the Biological Sciences Department

College of Science and Computer Studies

De La Salle University - Dasmariñas

Dasmariñas, Cavite

In Partial Fulfilment of the Requirementsfor the Degree

Bachelor of Science Major in Biology with Concentration in

Environmental Science

MA. PATRICIA B. ILAGAN

MARK ANTHONY P. BELTRAN

APRIL 2013

ABSTRACT

The aim of this study is to identify the different collected species of clams from Ang Pulo and correlate the amount of lead found in the clams to the amount of lead found in the water and benthic sediments in Ang Pulo Mangrove Reserve, Calatagan, Batangas. Clams are bivalves that are able to absorb heavy metals such as Lead in the environment by filtering the water. The collection of samples were done on the months of April, May and October to observe the difference in the amount of lead found in the samples during pre- wet and post- wet seasons.

Samples from Ang Pulo Mangrove Reserve, Calatagan, Batangas were collected by performing the convenient sampling method using quadrat method in three sampling stations once a month for three months. The clams were identified in the Institute of Environmental Science and Meteorology in the University of the Philippines Diliman. The samples underwent dry ashing and acid digestion before being subjected to the Atomic Absorption Spectrophotometer for the detection of Lead.

Seven (7) species of clams were collected from Ang Pulo Mangrove Reserve, Calatagan, Batangas. There is a very strong relationship between the amounts of lead found in the clams and the water samples. Similarly, the correlation between the amount of lead in the clams and the benthic sediments is very strong. The risk quotients of the clam and water samples were high while the risk quotient of the benthic sediments was low. The data also showed that there is a noticeable increase in the amount of lead found in the samples during the post wet season due to the accumulation of pollutants carried by the runoff from the wet season.



TABLE OF CONTENTS

Title Page
Approval Sheet
Acknowledgements
Abstract
Table of Contents
CHAPTER 1: INTRODUCTION
CIEM TEX I. II (INODOCTION)
1.1 Background of the study9
1.2 Conceptual framework1
1.3 General Objectives12
1.4 Scope and Limitations12
1.5 Significance of the study13
1.6 Definition of Terms14
CAHPTER 2: REVIEW OF RELATED LITERATURE
2.1 Conceptual Literature16
2.1 Conceptual Enterature
2.2 Related Studies
CHAPTER 3: METHODOLOGY
3.1 Research Design21
3.2 Research Setting21
3.3 Collection of Samples22
3.4 Collection of Data22
3.5 Analysis of Data23
CHAPTER 4: RESULTS AND DISCUSSION
4.1 Results
4.2 Discussion
CHAPTER 5: CONCLUSION AND RECOMMENDATION
5.1 Conclusion32
5.2 Recommendations33
GW 170 A
Cited References55

De La Salle University - Dasmariñas

Appendices	34
A. Map of Ang Pulo Calatagan, Batangas	
B. Atomic Absorption Spectrophotometer Preparation	
C. Computations and Statistics	37
D. Photodocumentation	
E. Certification of Identified Clams	46
F. Gantt Chart	53
G Curriculum Vitae	59





4.1 Lead Content of the collected Clams and their Risk Quotients		
4.2 Lead Content of Water in Calatagan, Batangas	25	
4.3 Lead Content of Benthic sediments in Calatagan, Batangas	26	





.1 Map of Ang Pulo Calatagan, Batangas	34
1.2 Preparation of sampling sites	43
1.3 Collection of Clams	43
1.4 Homogenization of Clams	44
1.5 Acid Digestion	44
1.6 Atomic Absorption Spectrophotometer	45
1.7 Clam Identification	45
1.8 Anadara antiquata	
1.9 Donax sp	47
1.10 Gafrarium pectinatum	48
1.11 Pitar citrinus	49
1.12 Tapes sp	50
1.13 Vasticardium enode	51
1.14 Vepricardium multispinosum	52