

# DE LA SALLE UNIVERSITY

TWO APPROACHES ON THE PROOF OF THE  
AMITSUR-LEVITZKI THEOREM

---

A Thesis  
Presented To  
the Faculty of Graduate School  
De La Salle University

In Partial Fulfillment  
of the Requirement for the Degree of  
Master of Science in Mathematics

---

by  
Bernardo L. Bagorio  
September 1991



# DE LA SALLE UNIVERSITY

## TABLE OF CONTENTS

	PAGE
ACKNOWLEDGMENT	0
CHAPTER	
I. THE PROBLEM AND ITS BACKGROUND	
1.1. Introduction	1
1.2. Statement of the Problem	1
1.3. Significance of the Study	2
1.4. Scope and Limitation	4
1.5. Methodology	4
1.6. Review of Related Literature	5
II. PRELIMINARY CONCEPTS	
2.1. Permutations	8
2.2. Algebraic Structures	21
2.3. Graph Theory	32
III. STANDARD POLYNOMIAL IDENTITY	40
IV. GRAPH THEORETIC PROOF OF AMITSUR- LEVITZKI THEOREM	66
V. SUMMARY, CONCLUSION AND RECOMMENDATIONS	83
BIBLIOGRAPHY	87



**CHAPTER V****SUMMARY**

We have presented here two methods of proving the Amitsur-Levitzki theorem. The graph theoretic proof was presented to give the readers the impression that there are several ways to prove a certain theorem. Inevitably, a reader would prefer a proof that is easy to understand, something that is less complicated. The simpler the proof is the better. The application of graph theory in proving Amitsur-Levitzki theorem has somehow minimized some complications. The employment of it has eliminated most of the complicated algebraic definitions merely by using much simpler graphic notions and by drawing a picture of the appropriate graph.

**CONCLUSION**

In this paper, we were able to prove the following theorems:



1. If  $F$  is a field, then  $M_m(F)$  satisfies the standard polynomial identity  $S_{2m}$ .

2. Let  $G$  be a graph of order  $n$ , let  $A$  be the set of arcs of  $G$  such that  $|A| = m$  and  $m \geq 2n$ . Let  $v$  and  $w$  be any fixed vertices of  $G$ . Then the number of unicursal paths  $\omega$  from  $v$  to  $w$  with  $\varepsilon(\omega) = +1$  is equal to the number of unicursal paths  $\omega$  from  $x$  to  $y$  with  $\varepsilon(\omega) = -1$ .

Aside from this, we were able to give some basic properties of polynomial identities and consequently the standard polynomial identities. Furthermore, the necessary and sufficient conditions for the existence of the unicursal paths in a given graph were discussed in this paper.

## RECOMMENDATIONS

Our recommendations in this paper, would focus on the algebraic aspect, particularly on polynomial identities. The concept of polynomial identities seems to suggest some special considerations for further studies. Hence, we recommend the following topics for others to do some research on.



1. We obtained in this study the result if  $E$  is an algebra over  $F$ , contained in  $M_m(F)$ , then it satisfies a polynomial identity, in fact a standard identity  $s_{2m}$ . The question here concerns the converse, that is, can all polynomial identity algebras be imbedded in suitable matrix rings? If so, what are those polynomial identity algebras that are imbeddable in matrix rings over commutative rings?

2. Another topic that might be of interest is the concept of generalized polynomial identities. Let  $E$  be an algebra over  $F$ . A generalized polynomial over  $E$  is, roughly speaking, a polynomial in the noncommuting indeterminates  $x_1, x_2, \dots, x_n$  in which the elements of  $E$  are allowed to appear both as coefficients and as entries between the indeterminates. Hence,  $E$  satisfies a generalized polynomial identity, if there exists a nonzero generalized polynomial  $f(x_1, x_2, \dots, x_n)$  such that

$$f(\alpha_1, \alpha_2, \dots, \alpha_n) = 0$$

for all  $\alpha_i$ 's  $\in E$ . The problem here is to find the conditions that will ensure the existence of a



generalized polynomial identity.

