



**SCREENING AND IDENTIFICATION OF THE ANTIBIOTIC-
PRODUCING SOIL ACTINOMYCETES FROM A DUMPSITE IN LOS
BAÑOS, LAGUNA AGAINST A PANEL OF PATHOGENIC BACTERIA**

An Undergraduate Research Presented to the
Faculty of the Biological Sciences Department
College of Science
De La Salle University - Dasmariñas
Dasmariñas, Cavite

In Partial Fulfilment of the Requirements
for the Degree of Bachelor of Science Major in Human Biology

KATHERINE L. PINTOR

March 2012



ABSTRACT

The rapid emergence of drug resistance among various pathogenic bacteria stresses the need to look for new compounds used in the production of antibiotics. This study aimed to isolate novel actinomycetes that can be a new source of inhibitory compounds to various pathogenic organisms. The soil samples were collected from a dumpsite in Los Baños, Laguna. The actinomycetes were isolated in specific medium using Yeast-Malt extract Agar (YMA). A total of 44 Actinomycetes strains were isolated and screened for their *in vitro* antagonistic activities against a panel of test organisms which were: Methicillin-resistant *Staphylococcus aureus* (MRSA), *Klebsiella pneumoniae* (KP), *Staphylococcus epidermidis* (SE), *Pasteurella multocida* (PM), *Staphylococcus aureus* (SA), and *Pseudomonas aeruginosa* (PA). The actinomycetes strains were subjected to preliminary screening using agar plug method to determine their inhibitory activity. The results indicated that a broad spectrum of inhibitory activity was observed in some of the isolates. These strains can be potential source for new compounds for antibiotic production against different pathogens. Moreover, the six most potent actinomycetes strains with the largest broad spectrum of antibacterial activity against each of the test organisms were selected. The isolates were then subjected to cup cylinder assay. The isolates have shown ability to produce substances that can inhibit drug resistant bacteria. In addition, the best isolate which displayed the greatest antagonistic activity in the cup cylinder assay against any of the test organisms was selected and identified. The isolate A20 was identified to be *Tsukamurella tyrosinosolvans* M12400 using 16S ribosomal RNA gene sequencing method. This study signifies that actinomycetes isolated in extreme environment such as landfills can be a great source of bioactive compounds.



TABLE OF CONTENTS

Title Page	1
Table of Contents	6
CHAPTER 1 INTRODUCTION	
1.1 Background of the Study	8
1.2 Conceptual Framework	10
1.3 Statement of the Problem	12
1.4 Scope and Limitations	13
1.5 Significance of the Study	14
1.6 Definition of Terms	16
CHAPTER 2 LITERATURE REVIEW	
2.1 Conceptual Literature	18
2.2 Related Studies	33
CHAPTER 3 METHODOLOGY	
3.1 Research Design	44
3.2 Research Setting	44
3.3 Research Procedure	44
3.4 Data Gathering	48
CHAPTER 4 RESULTS AND DISCUSSION	
4.1 Results	49
4.2 Discussion	54



CHAPTER 5 CONCLUSION AND RECOMMENDATION

5.1 Conclusion	60
5.2 Recommendation	61
Cited References	63
Appendices	
A. Map of Study Site	75
B. Standard Procedure	76
C. Results of 16S ribosomal RNA gene sequencing	89
D. Extraction of Genomic DNA	92
E. PCR Amplification	94
F. Restriction Endonuclease Digestions & Nuclease	95
G. Gantt Chart	96
H. Budget for Experimentation	97
Curriculum Vitae	98